



Site Marked- Left / Right

Consent Timeout performed

Indication

Surgical Anesthesia Surgeon request for post-operative pain management

Nerve Block Procedure

Interscalene Supraclavicular Infraclavicular Axillary
 Lumbar Plexus Femoral Sciatic/Popliteal Ankle
 TAP Block- Bilateral
 IPACK

Ultrasound Guided- I confirm that real time ultrasound was utilized to to guide the needle for satisfactory placement and Ultrasound guidance was used to view medication diffusion. Image was printed and attached to record.

Monitors- Blood Pressure Pulse Oximetry EKG ETCO2

Oxygen _____ (L/min)

Nasal Cannula FaceMask

Premedications

Midazolam _____ mg Fentanyl _____ mcg Propofol _____ mg Other _____

Level of Sedation

Awake Sedated/Easily aroused General Anesthesia Spinal Anesthesia

Needle

2 inch 4 inch Other _____

Local Anesthetic

Type	Concentration	Volume
<input type="checkbox"/> Bupivacaine		
<input type="checkbox"/> Ropivacaine		
<input type="checkbox"/> Lidocaine		

Procedure Note

Start time _____ End Time _____

Position Supine Lateral Prone

Patient prepped with antiseptic

Nerve Stimulator minimal current _____ mA not used

Blood Aspirated No Yes-Action taken _____

Paresthesia No Yes-Action taken _____

Pain on injection No Yes-Action taken _____

Signature _____

Date/Time _____